


HOMOGENIZATION IN RANDOM MEDIA AND ISOTROPY CONSIDERATIONS

Alen Alexanderian, Muruhan Rathinam, and Rouben Rostamian
University of Maryland, Baltimore County

DIFFUSION IN A RANDOM MEDIUM




$$\begin{cases} -\operatorname{div}_{\mathbf{x}} (A(\mathbf{x}, \omega) \nabla u(\mathbf{x}, \omega)) = f(\mathbf{x}) & \text{on } \mathcal{D} \\ u(\mathbf{x}, \omega) = 0 & \text{on } \partial\mathcal{D} \end{cases}$$

Some considerations:

- Random PDE \Rightarrow difficult problem
- For fixed realization computation is a formidable task
- Need to have a discretization of the domain fine enough to resolve the complicated micro-structure
- The problem is either too computationally expensive or impossible to solve numerically

HOMOGENIZATION

Physical intuition: instead of working with a medium with complicated structure work with a homogeneous medium with the same effective properties.



Consider the problem:

$$\begin{cases} -\operatorname{div}_{\mathbf{x}} (A(\mathbf{x}, \omega) \nabla u(\mathbf{x}, \omega)) = f(\mathbf{x}) & \text{on } \mathcal{D} \\ u(\mathbf{x}, \omega) = 0 & \text{on } \partial\mathcal{D} \end{cases} \quad (1)$$


Given appropriate conditions on A , there is a homogenized problem [4, 2]:

$$\begin{cases} -\operatorname{div}(A^0 \nabla u^0) = f & \text{on } \mathcal{D} \\ u^0 = 0 & \text{on } \partial\mathcal{D} \end{cases} \quad (2)$$

such that for almost all ω , the solution u^0 of (2) is a close approximation of the solution u of the problem (1).

SYMMETRIES OF RANDOM MEDIA

- Let Q be a rotation matrix
- A random structure is invariant under rotation by Q if rotation of a realization by Q gives an equally likely realization



ISOTROPY (DIFFUSION)

- A medium is isotropic if it behaves the same regardless of its orientation
- Consider the homogenized problem


$$\begin{cases} -\operatorname{div}(A^0 \nabla u^0) = f & \text{on } \mathcal{D} \\ u^0 = 0 & \text{on } \partial\mathcal{D} \end{cases} \quad (3)$$

$$\text{Isotropy} \Leftrightarrow A^0 = \alpha I.$$

Question: Under what conditions the effective behavior of a composite medium is isotropic?


Answer in 2D:

- R_θ matrix of rotation by angle θ
- If $\theta \neq k\pi$, invariance under $R_\theta \Rightarrow$ isotropy



NUMERICAL COMPUTATIONS

Consider a random checkerboard:




- Random checkerboard, fifty-fifty chance of white or gray
- gray conductivity $a_1 = 1$
- white conductivity $a_2 = 2$

We expect [2],

$$A^0 = \sqrt{a_1 a_2} I = \sqrt{2} I$$

PERIODIZATION



- Fix a realization, select a square $[0, \rho]^2$, and extend it periodically to \mathbf{R}^2
- Solve the unit cell problem from periodic homogenization
- Get the corresponding homogenized matrix $A^\rho(\omega)$

Then [1, 3],

$$\text{As } \rho \rightarrow \infty, A^\rho \rightarrow A^0 \text{ for almost all } \omega.$$

SAMPLE COMPUTATIONAL RESULTS

$$A_1^\rho = \begin{bmatrix} 1.2397 & -0.0029 \\ -0.0029 & 1.2580 \end{bmatrix}$$

$$A_2^\rho = \begin{bmatrix} 1.3876 & 0.0013 \\ 0.0013 & 1.4074 \end{bmatrix}$$

$$A_3^\rho = \begin{bmatrix} 1.4099 & 0.0015 \\ 0.0015 & 1.4101 \end{bmatrix}$$

LINEAR ELASTICITY

- \mathcal{E} : n -dimensional Euclidean space
- $\mathcal{D} \subset \mathcal{E}$

$$\begin{cases} -\operatorname{div} C[\nabla \mathbf{u}] = \mathbf{f}, & \text{in } \mathcal{D}, \\ \mathbf{u} = 0 & \text{on } \partial\mathcal{D}. \end{cases}$$

- $\mathbf{f} \in (L^2(\mathcal{D}))^n$: body force
- C : elasticity tensor

$$C : \operatorname{Sym}(\mathcal{E}) \rightarrow \operatorname{Sym}(\mathcal{E})$$

ISOTROPY (ELASTICITY)

Definition: Symmetry group of an elasticity tensor C is a group $G_c \subseteq \operatorname{Orth}(\mathcal{E})$ such that

$$QC[E]Q^T = C[QEQ^T], \forall Q \in G_c. \quad (4)$$

C is Isotropic if

$$QC[E]Q^T = C[QEQ^T], \quad \forall Q \in \operatorname{Orth}(\mathcal{E}). \quad (5)$$

Question: Under what condition on G_c does (4) \Rightarrow (5)?

Answer in 2D:

- \mathcal{E} : the 2-dimensional Euclidean space
- $R_\theta \in \operatorname{Orth}(\mathcal{E})$: rotation by angle θ
- If $\theta \neq k\frac{\pi}{2}$, invariance under $R_\theta \Rightarrow$ isotropy

REFERENCES

- [1] A. Bourgeat and A. Piatnitski. Approximations of effective coefficients in stochastic homogenization. *Annales de l'Institut Henri Poincaré (B) Probability and Statistics*, 40 (2), 2004.
- [2] V. V. Jikov, S. M. Kozlov, and O. A. Olenik. *Homogenization of differential operators and integral functionals*. Springer, 1994.
- [3] H. Owhadi. Approximation of the effective conductivity of ergodic media by periodization. *Probab. Theory Related Fields*, 125(2), 2003.
- [4] G. C. Papanicolaou and S. R. S. Varadhan. Boundary value problems with rapidly oscillating random coefficients. *Seria Colloquia Mathematica Societatis Janos Bolyai North-Holland*, 27, 1981.